

$$(1) \quad F(t, y(t), y'(t)) = 0$$

$$F: \mathbb{R}^3 \rightarrow \mathbb{R}$$

$$(2) \quad y' = f(t, y(t))$$

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}.$$

$$F(\alpha, \beta, y) = \alpha + \beta + y$$

$$t + y(t) + y'(t) = 0$$

$$y'(t) = -t - y(t)$$

$$y'(t) = f(t, y(t))$$

$$f(\alpha, \beta) = -\alpha - \beta.$$

$$F(t, y, y') = 0 \quad y' = f(t, y)$$

$$y'' = y - y'.$$

$$F(t, y, y') = 0 \quad \varphi: \mathbb{R} \rightarrow \mathbb{R} \quad D\varphi = J$$

$$\forall t \in J : \quad F(t, \varphi(t), \varphi'(t)) = 0 \quad \Rightarrow$$

φ je řešením na J .

Pr. $\varphi(t) = 1-t$ je na $J = (-\infty, \infty)$ řešením

$$\text{ODR} \text{ a vyu} \quad t + y + y' = 0.$$

$$\forall t \in (-\infty, \infty) :$$

$$t + \psi(t) + \psi'(t) = t + (1-t) + (-1) = 0 .$$

$$\psi(t) = e^{-t} + (1-t)$$

$$\forall t \in (-\infty, \infty) :$$

$$t + \psi(t) + \psi'(t) = t + e^{-t} + (1-t) + (-e^{-t} - 1) = \\ = 0 .$$

$$t + y + y' = 0$$

$$y(t) = C \cdot e^{-t} + (1-t), \quad C \in \mathbb{R} .$$

$$(y')^2 + 1 = 0 \quad \forall t \in J : (y'(t))^2 = -1 .$$

$$\underline{F(t, y, y')} = 0 \quad \underline{y(t_0) = y_0} .$$

Cauchyova úloha

$$t + y + y' = 0 , \quad y(0) = 2$$

$$\psi(t) = e^{-t} + (1-t) , \quad \psi(0) = 2 .$$

$$\underline{F(t, y, y')} = 0 \quad \begin{matrix} \varphi_1 & \text{na} & J_1 \\ \varphi & \text{na} & J \end{matrix}$$

$$J_1 \subset J, \quad J_1 \neq J$$

$$\forall t \in J_1 : \quad \psi_1(t) = \psi(t) \quad \varphi \text{ je prohlouzením } \varphi_1 \text{ na } J$$